CONTENTS

(Abstracts/contents lists published in: Chem. Abstr.; Curr. Contents, Phys. Chem. Earth Sci., Life Sci.)

Introduction ix

An asymmetric approach to 2-deoxynucleosides *via* organosulfur building blocks as chemical chameleons
B. M. Trost and C. Nübling (Stanford, CA, U.S.A.) 1

De novo synthesis of carbohydrates by stereoselective aldol reaction: L-cladinose
S. H. Montgomery, M. C. Pirrung, and C. H. Heathcock (Berkeley, CA, U.S.A.) 13

Synthesis of L-gulose, L-galactose, and their acetylated *aldehydo* forms from 6-*S*-phenyl-6-thio-D-hexoses
F. Santoyo González and H. H. Baer (Ottawa, Ont., Canada) 33

Multiple and long-range participation of benzyl groups in intramolecular *C*-arylation reactions of benzylated glycosides
O. R. Martin, S. P. Rao, C. A. V. Hendricks, and R. E. Mahnken (Binghampton, NY, U.S.A.) 49

Novel methods for the preparation of partially acetylated carbohydrates
S. Hanessian and M. Kagotani (Montreal, Que., Canada) 67

1,6-Anhydro-β-D-glucopyranose in organic synthesis: preparation of a fragment for the synthesis of rosaramycin
G. Procter, D. Genin (Salford, Gt. Britain), and S. Challenger (Cardiff, Gt. Britain) . . . 81

Inversion of configurations of contiguous carbinol centers: application to the synthesis of both enantiomers of natural products from the same enantiomerically pure starting material
P. Prasit, G. Robertson, and J. Rokach (Pointe Claire-Dorval, Que., Canada) 93

Effect on human liver glycosidases and short syntheses of 1α,2α,6α,7aβ-1,2,6,7-tetrahydroxypyrrolizidine from D-*glycero*-D-*gulo*-heptono-1,4-lactone
W. F. Collin, G. W. J. Fleet, M. Haraldsson (Oxford, Gt. Britain), I. Cenci di Bello, and B. Winchester (London, Gt. Britain) 105

Synthesis of 1-deoxy-6-epicastanospermine and 1-deoxy-6,8a-diepicastanospermine
K. H. Aamlid, L. Hough, and A. C. Richardson (London, Gt. Britain) 117

Derivatives of methyl β-lactosides as substrates for and inhibitors of β-D-galactosidase from *E. coli*
K. Bock (Copenhagen, Denmark), and K. Adelhorst (Lyngby, Denmark) 131

The use of *N*-alkoxycarbonyl derivatives of 2-amino-2-deoxy-D-glucose as donors in glycosylation reactions
P. Boullanger, M. Jouineau, B. Bouammali, D. Lafont, and G. Descotes (Villeurbanne, France) 151

Benzeneselenenyl triflate as an activator of thioglycosides for glycosylation reactions
Y. Ito, T. Ogawa (Wako, Japan), M. Numata, and M. Sugimoto (Tokorozawa, Japan) . . 165

Total synthesis of globotriaosylceramide (Gb_3) and lysoglobotriaosylceramide (lysoGb$_3$)
K. C. Nicolaou, T. J. Caulfield, and H. F. Katoaka (La Jolla, CA, U.S.A.) 177

Muramic acid derivatives as glycosyl donors for the synthesis of muramyl-containing glycosphingolipids and fatty acids
A. Toepfer and R. R. Schmidt (Konstanz, F.R.G.) 193

Synthesis of glycopeptides with the T_N and T antigen structures, and their coupling to bovine serum albumin
H. Kunz, S. Birnbach, and P. Wernig (Mainz, F.R.G.) 207

Use of the methylsulfenyl cation as an activator for glycosylation reactions with alkyl (aryl) 1-thioglycopyranosides: synthesis of methyl *O*-(2-acetamido-2-deoxy-*β*-D-glucopyranosyl)-(1→6)-*O*-*α*-D-glucopyranosyl-(1→2)-*α*-D-glucopyranoside, a derivative of the core trisaccharide of *E. coli* K 12
F. Dasgupta and P. J. Garegg (Stockholm, Sweden) 225

1,3,4,6-Tetra-*O*-acetyl-2-chloroacetamido-2-deoxy-*β*-D-glucopyranose as a glycosyl donor in synthesis of oligosaccharides
F. Dasgupta and L. Anderson (Madison, WI, U.S.A.) 239

Synthesis of trisaccharide C–D–E and tetrasaccharide B–C–D–E fragments found in orthosomysins
M. Trumtel, P. Tavecchia, A. Veyrières, and P. Sinaÿ (Paris, France) 257

Author index c1

Subject index c3